

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ended August 5 there occurred 17 cases and 13 deaths, viz: 15 cases, 12 deaths, in Alexandria; 1 case, 1 death, in Dekernes, and 1 case in Port Said.

Death rate of Berlin and other cities.

The death rate of Berlin for the week ended August 5 was higher than it has been at almost any time during the past 12 months, amounting, calculated on the year, to 21.4 per thousand of the population (against 17.8 in the preceding week), this being, however, a little lower than the rate for the same week of last year, in which it amounted to 22.5 per thousand. Notwithstanding the increased mortality in Berlin, two-thirds of the large towns and cities of Germany showed more unfavorable health conditions than this city, the following cities having a considerably higher death rate than Berlin, namely: Nuremberg, Stuttgart, Frankfort on the Main, Bremen, Brunswick, Dresden, Rixdorf (with 32.8), Königsberg, Breslau, Halle, Leipzig, Magdeburg, Cologne, Karlsruhe, Strassburg. The following towns showed a lower rate of mortality than Berlin, viz: Hamburg, Altona, Munich, Cassel, Hanover, Schöneberg (with 14 per thousand), Charlottenburg (with the minimum of 13.3); also London, Paris, and Vienna. foregoing week the increase of the mortality was exclusively among children in the first year of life, there being, indeed, a decrease in the mortality among the higher age classes. The infant death rate rose from 7.2 per year and thousand in the preceding week to 11.1 in this week, thus corresponding with the death rate of the same week of last year; it was higher than the Hamburg and Munich rate, but lower than the Leipzig figure. There were registered 330 deaths from catarrh of the stomach and intestines (307 infants and 23 adults), 106 deaths from phthisis pulmonalis, 61 deaths from diseases of the respiratory organs, 53 deaths from cancer, 16 deaths from measles, 8 deaths from diphtheria, 4 deaths from scarlet fever, 1 death from enteric fever. Finally, 3 persons died by violence.

GUATEMALA.

Report from Livingston, fruit port.

Acting Assistant Surgeon Peters reports as follows: Week ended August 19, 1905. Present officially estimated population, 3,500; no deaths; prevailing disease, malarial fever of mild type; general sanitary condition of this port and the surrounding country during the week, fair. The last death from yellow fever occurred on July 30, since which date no cases have come under observation. The health of Livingston and Puerto Barrios at present is good. There is very little sickness in either place. No deaths have occurred during the week.

Bills of health were issued to the following-named vessels:

Date.	Vessel.	Destination.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of bag gage disin- fected.
Aug. 15	Olympia	Mobile via	42 24			
		Belize.				